

## Abstract

We analyse the relationship between different extremal notions in Lipschitz free spaces. We completely characterise strongly exposed points in the unit ball of a free space. We prove that every preserved extreme point of the unit ball is also a denting point. We show in some particular cases that every extreme point is a molecule, and that a molecule is extreme whenever the two points, say  $x$  and  $y$ , which define it satisfy that the metric segment  $[x, y]$  only contains  $x$  and  $y$ . As an application, we get some new consequences about norm-attainment in spaces of vector-valued Lipschitz functions. This is based in a joint work with A. Procházka, C. Petitjean and A. Rueda Zoca.

## Introduction


We are interested in studying the following families of distinguished points in the unit ball  $B_X$  of a Banach space  $X$ .

### Definition

A point  $x$  in the unit ball  $B_X$  of a Banach space  $X$  is said to be:

- an **extreme point** of  $B_X$  if  $x = \lambda y + (1 - \lambda)z$ ,  $y, z \in B_X$ ,  $\lambda \in (0, 1)$ , then  $x = y = z$ .
- a **preserved extreme point** of  $B_X$  if  $x$  is an extreme point of  $B_{X^{**}}$ ,
- a **denting point** of  $B_X$  if the slices of  $B_X$  containing  $x$  are a neighbourhood basis of  $x$  in  $(B_X, \|\cdot\|)$ .
- an **exposed point** of  $B_X$  if there is  $f \in X^*$  such that  $f(x) > f(y)$  for every  $y \in B_X \setminus \{x\}$ .
- a **weak-strongly exposed point** of  $B_X$  if there is  $f \in X^*$  such that for every sequence  $(x_n)_n$  in  $B_X$  we have  $x_n \xrightarrow{w} x$  whenever  $f(x_n) \rightarrow f(x)$ , equivalently, the slices of  $B_X$  provided by  $f$  are a neighbourhood basis of  $x$  in  $(B_X, w)$ .
- a **strongly exposed point** of  $B_X$  if there is  $f \in X^*$  such that for every sequence  $(x_n)_n$  in  $B_X$  we have  $x_n \rightarrow x$  whenever  $f(x_n) \rightarrow f(x)$ , equivalently, the slices of  $B_X$  provided by  $f$  are a neighbourhood basis of  $x$  in  $(B_X, \|\cdot\|)$ .

It is not difficult to check that the above concepts are related in the following way:



Moreover, none of these implications reverse in general.

Our aim is to study the former notions in the particular case in which the Banach space is the Lipschitz free space  $\mathcal{F}(M)$  over a metric space  $(M, d)$ . Recall that the space  $\text{Lip}_0(M)$  of Lipschitz functions on  $M$  vanishing at a distinguished point  $0 \in M$  is a Banach space when it is endowed with the norm given by the best Lipschitz constant. Then

$$\mathcal{F}(M) := \overline{\text{span}}\{\delta(x) : x \in M\} \subset \text{Lip}_0(M)^*,$$

where  $\langle \delta(x), f \rangle = f(x)$  for  $f \in \text{Lip}_0(M)$ . We refer the reader to [1, 2] for the fundamental properties and applications of Lipschitz free spaces. Let us highlight that for every Banach space  $Y$  and every Lipschitz function  $f : M \rightarrow Y$  such that  $f(0) = 0$  there is a unique bounded linear operator  $\hat{f} : \mathcal{F}(M) \rightarrow Y$  such that  $\hat{f} \circ \delta = f$ . Moreover,  $\|\hat{f}\| = \|f\|$ . It follows from this fact that  $\mathcal{F}(M)^*$  is isometric to  $\text{Lip}_0(M)$ .

The study of the extremal structure of  $B_{\mathcal{F}(M)}$  probably was started by Weaver in [2], where it is proved that **every preserved extreme point of  $B_{\mathcal{F}(M)}$  is a molecule**, that is, an element of the form

$$m_{xy} = \frac{\delta(x) - \delta(y)}{d(x, y)}, \quad x, y \in M, x \neq y.$$

We denote  $V$  the set of molecules in  $\mathcal{F}(M)$ . Note that

$$\|f\| = \sup \left\{ \frac{f(x) - f(y)}{d(x, y)} : x, y \in M, x \neq y \right\} = \sup \{ \langle f, m_{xy} \rangle : m_{xy} \in V \}$$

and so  $V$  is 1-norming for  $\text{Lip}_0(M)$ . Equivalently,


$$B_{\mathcal{F}(M)} = \overline{\text{conv}}(V).$$

That provides a useful way for describing the norm in  $\mathcal{F}(M)$ .

## Extreme points in $B_{\mathcal{F}(M)}$

In order to get some intuition, let us assume first that  $M$  just consists of three points,  $M = \{0, x, y\}$ . Let us define the following metrics on  $M$ :

$$d_1(x, 0) = d_1(y, 0) = d_1(x, y) = 1 \quad d_2(x, 0) = d_2(y, 0) = 1, d_2(x, y) = 2$$



Note that in  $\mathcal{F}(M, d_1)$  the set of extreme points of the ball coincides with the set of molecules. On the other hand, the molecule  $m_{xy}$  is not an extreme point of the ball of  $\mathcal{F}(M, d_2)$ . The reason is that  $0$  belongs to the metric segment  $[x, y]$  in  $(M, d_2)$ . More generally, for a molecule  $m_{xy}$  to be an extreme point of  $B_{\mathcal{F}(M)}$  it is necessary that the metric segment  $[x, y]$  between  $x$  and  $y$  reduces to  $\{x, y\}$ . Indeed, if  $d(x, z) + d(z, y) = d(x, y)$  for some  $z \in M \setminus \{x, y\}$  then

$$m_{xy} = \frac{d(x, z)}{d(x, y)} m_{xz} + \frac{d(z, y)}{d(x, y)} m_{zy}$$

and so  $m_{xy}$  is not an extreme point of  $B_{\mathcal{F}(M)}$ . This fact motivates the following question:

### Open problem

Assume that  $[x, y] = \{x, y\}$ . Is  $m_{xy}$  an extreme point of  $B_{\mathcal{F}(M)}$ ?

Aliaga and Guirao have recently proved that the above problem has an affirmative answer if  $M$  is compact. We have shown the following:

Let  $M$  be a bounded uniformly discrete (i.e.  $\inf_{x \neq y} d(x, y) > 0$ ) metric space. If  $[x, y] = \{x, y\}$ , then  $m_{xy}$  is an extreme point of  $B_{\mathcal{F}(M)}$ .

The above result allow us to find an example of a bounded uniformly discrete countable metric space  $M$  such that  $\mathcal{F}(M)$  is not isometric to a dual Banach space.

Another natural question is the following:

### Open problem

If  $\mu$  is an extreme point of  $B_{\mathcal{F}(M)}$ , is  $\mu$  necessarily a molecule  $\mu = m_{xy}$ ?

We have shown that this question has an affirmative answer in some particular cases.

### Theorem 1 (GL – Procházka – Petitjean – Rueda Zoca, 2017)

Let  $M$  be a bounded separable metric space. Assume that there is a subspace of  $\text{lip}_0(M)$  (little-Lipschitz functions) which is predual of  $\mathcal{F}(M)$  and  $\delta(M)$  is weak\*-closed. Then given  $\mu \in B_{\mathcal{F}(M)}$  the following are equivalent:

- $\mu$  is an extreme point of  $B_{\mathcal{F}(M)}$ .
- $\mu$  is an exposed point of  $B_{\mathcal{F}(M)}$ .
- There are  $x, y \in M$ ,  $x \neq y$ , such that  $[x, y] = \{x, y\}$  and  $\mu = m_{xy}$ .

This applies in the following cases:

- $M$  compact countable.
- $(M, d^\alpha)$ ,  $0 < \alpha < 1$  compact  $\alpha$ -snowflaking of a metric space  $(M, d)$ .
- $M$  bounded uniformly discrete admitting a compact topology  $\tau$  such that  $d$  is  $\tau$ -lsc.

Theorem 1 has the following application to the norm-attainment of Lipschitz functions, which extends a result in [1].

Let  $Y$  be a Banach space and  $M$  be a metric space satisfying the hypotheses of Theorem 1. Then every Lipschitz function  $f : M \rightarrow Y$  which attains its norm as an operator from  $\mathcal{F}(M)$  to  $Y$  also attains its Lipschitz norm on a pair of points in  $M$ .

## Strongly exposed points in $B_{\mathcal{F}(M)}$

Weaver proved that  $m_{xy}$  is a preserved extreme point of  $B_{\mathcal{F}(M)}$  whenever there is  $f \in \text{Lip}_0(M)$  peaking at  $(x, y)$ , that is,  $\langle f, m_{xy} \rangle = 1$  and  $\sup_{(u,v) \notin U} \langle f, m_{uv} \rangle < 1$  for every open subset  $U$  of  $M^2 \setminus \Delta$  containing  $(x, y)$  and  $(y, x)$ . Indeed, peaking functions characterise strongly exposed points in  $B_{\mathcal{F}(M)}$ .

### Theorem 2 (GL – Procházka – Rueda Zoca, 2017)

Let  $x, y \in M$ ,  $x \neq y$ . The following assertions are equivalent:

- The molecule  $m_{xy}$  is a strongly exposed point of  $B_{\mathcal{F}(M)}$ .
- There is  $f \in \text{Lip}_0(M)$  peaking at  $(x, y)$ .
- There is  $\varepsilon > 0$  such that  $d(x, z) + d(z, y) - d(x, y) > \varepsilon \min\{d(x, z), d(z, y)\}$  for all  $z \in M \setminus \{x, y\}$ .

This result extends a characterisation of peaking functions in subsets of  $\mathbb{R}$ -trees due to Dalet, Kaufmann and Procházka [3].

It was shown in [4] that if  $M$  is compact then  $\text{Lip}_0(M)$  has the Daugavet property if and only if condition (iii) fails for every pair of distinct points in  $M$ . As a consequence, the following dichotomy holds:

Let  $M$  be a compact metric space. Then either  $\text{Lip}_0(M)$  has the Daugavet property (and so every slice of  $B_{\mathcal{F}(M)}$  has diameter 2) or  $B_{\mathcal{F}(M)}$  has a strongly exposed point.

## Preserved extreme points in $B_{\mathcal{F}(M)}$

Aliaga and Guirao have recently proved a characterisation of preserved extreme points in  $B_{\mathcal{F}(M)}$  in the spirit of Theorem 2. Namely, they prove in [5] that  $m_{xy}$  is a preserved extreme point of  $B_{\mathcal{F}(M)}$  if and only if for every  $\varepsilon > 0$  there is  $\delta > 0$  such that

$$(1 - \delta)(d(x, z) + d(z, y)) < d(x, y), z \in M \setminus \{x, y\} \implies \min\{d(x, z), d(y, z)\} < \varepsilon.$$

We have proved the following result:

### Theorem 3 (GL – Procházka – Petitjean – Rueda Zoca, 2017)

Every preserved extreme point of  $B_{\mathcal{F}(M)}$  is a denting point, and every weak-strongly exposed point of  $B_{\mathcal{F}(M)}$  is a strongly exposed point.

Now, one may wonder if some more implications in the diagram hold in the particular case of  $B_{\mathcal{F}(M)}$ . However, we have shown:

- There is a compact metric space  $M$  with a denting point of  $B_{\mathcal{F}(M)}$  which is not strongly exposed.
- There is a uniformly discrete countable metric space  $M$  with an exposed point of  $B_{\mathcal{F}(M)}$  which is not a preserved extreme point.

Finally, a curious consequence of Theorem 3 is the following:

The norm of  $\text{Lip}_0(M)$  is Gâteaux differentiable at  $f$  if and only if it is Fréchet differentiable at  $f$ .

## References

- [1] G. Godefroy, "A survey on Lipschitz-free Banach spaces," *Comment. Math.*, vol. 55, no. 2, pp. 89–118, 2015.
  - [2] N. Weaver, *Lipschitz algebras*. World Scientific Publishing Co., Inc., River Edge, NJ, 1999.
  - [3] A. Dalet, P. L. Kaufmann, and A. Procházka, "Characterization of metric spaces whose free space is isometric to  $\ell_1$ ," *Bull. Belg. Math. Soc. Simon Stevin*, vol. 23, no. 3, pp. 391–400, 2016.
  - [4] Y. Ivakhno, V. Kadets, and D. Werner, "The Daugavet property for spaces of Lipschitz functions," *Math. Scand.*, vol. 101, no. 2, pp. 261–279, 2007.
  - [5] R. Aliaga and A. J. Guirao, "On the preserved extremal structure of Lipschitz-free spaces." arXiv:1705.09579, 2017.
  - [6] L. García-Lirola, A. Procházka, and A. Rueda Zoca, "A characterisation of the Daugavet property in spaces of Lipschitz functions." arXiv:1705.05145, 2017.
  - [7] L. García-Lirola, C. Petitjean, A. Procházka, and A. Rueda Zoca, "Extreme structure and duality in Lipschitz free spaces." arXiv:1705.05145, 2017.
- The research of L. García-Lirola was supported by the grants MINECO/FEDER MTM2014-57838-C2-1-P and Fundación Séneca CARM 19368/PI/14.